Recovery Kernel

DISCONTINUANCE PROTOCOL: GHB / GBL / 1,4-BD

Die Vinnigste, Mees Onvoorspelbare Terugval in Hierdie Serie – en Hoe om Dit Veilig Te Oorleef

---

Every other guide in this series has a point where it says 'this is uncomfortable but rarely dangerous.' This guide doesn't get to say that. GHB withdrawal, in anyone dosing more than three or four times daily, is a genuine medical emergency with a real risk of seizure, a specific and severe delirium, and death without proper management. This isn't dramatized for effect — it's the actual, documented clinical picture, and treating it with anything less than full seriousness is how people get hurt. This guide exists to make sure the seriousness translates into a plan, not just fear.

Section 0 — Immediate Safety: Read This First

MEDICAL EMERGENCY

AS YOU ARE DOSING GHB, GBL, OR 1,4-BD MORE THAN THREE OR FOUR TIMES DAILY, DO NOT STOP WITHOUT MEDICAL SUPERVISION. WITHDRAWAL CAN BEGIN WITHIN 2-6 HOURS OF THE LAST DOSE, AND THE FIRST 12-72 HOURS CARRY A REAL, UNPREDICTABLE RISK OF SEIZURE AND A SEVERE DELIRIUM/PSYCHOSIS SYNDROME THAT CAN BE FATAL WITHOUT PROPER MEDICAL MANAGEMENT. GO TO AN EMERGENCY DEPARTMENT NOW IF YOU ARE A DEPENDENT USER PLANNING TO STOP, OR IF WITHDRAWAL HAS ALREADY BEGUN.

Signs that you are already in dangerous withdrawal — go to an ER now:

- Enige krampsaktiwiteit, hoe kort ook
- Verwarring, desoriëntasie of halusinasies – dit kan vinnig ontwikkel en is nie dieselfde as gewone angs nie.
- Ekstreme agitering gekombineer met 'n rennende hart, hoë bloeddruk, of swaar sweet (outonoom instabiliteit)
- A tremor that is getting worse rather than staying steady

⚠ WAARSKEUWING

There is no reversal agent for GHB overdose and no single FDA-approved medication for GHB withdrawal — unlike opioids (Naloxone) or alcohol (benzodiazepines have a clear, standard protocol). This means the quality of care depends heavily on the treating physician's familiarity with GHB specifically. Section 8 of this guide exists to help you or someone supporting you advocate for the right treatment if the emergency team seems unfamiliar with it.

Afdeling 1 — Hoekom Hierdie Onttrekking Verskil Van Elke Ander Gids in Hierdie Serie

What Makes It Different	Hoekom Dit Saakmaak
Onset speed	GHB se plasma half-life is net 30-60 minute. Withdrawal begin ek spoel van die laaste dosis by 'n afhanklike gebruiker — baie vinniger as alkohol (6-24 uur) of benzodiazepines (12-48 uur). Daar is baie min waarskuwingvenster tussen 'laaste dosis' en 'gevaar.'
Unpredictability	Klinieke ernsheid volg nie altyd skoon met gerapporteerde gebruik nie – iemand wat hul gebruik onderrapporteer, of wie se voorraad onstabiele konsentrasie gehad het (uiters algemeen met GHB-prekursore soos GBL en 1,4-BD), kan baie ernstiger voorgestel word as verwag.
'n aparte delirium en psigose syndroom	This ain't normal anxiety building up. GHB withdrawal delirium involves being jumpy, confused, seeing things that aren't there, and wild swings in your body's automatic systems—a real neurological emergency, different from the way booze causes delirium tremens.
No standaard behandelingsprotokol	Alcohol withdrawal has a clear, widely-known benzodiazepine protocol. Opioid withdrawal has buprenorphine and methadone. GHB withdrawal has no FDA-approved specific treatment — baclofen (a GABA-B agonist, the same receptor system GHB acts on) is the most evidence-supported option, but many emergency physicians have limited direct experience with it.

Section 2 — Die Ontwenning Tydlyn

Timeline	What's happening?	What to Expect	What to Do
Ureko 2-6 (Begin — die mediese ingrypingsvenster)	GHB clearing rapidly; GABA-B receptors left without their ligand; glutamate surging; HPA axis activating	Nervousness, tremors, sweating, a racing heart, queasiness — the start is quick and can be confused with normal worry until it builds up rapidly.	Dit is wanneer mediese ingryping (baclofen of benzodiazepines, Afdeling 8) moet begin as 'n afhanklike gebruiker stop. Indien nie reeds in mediese sorg is en hierdie simptome verskyn: gaan nou na die ER.
Hours 12-72 (Peak danger window)	Peak GABA-B receptor withdrawal; die hoogste-risiko venster vir krampe en die	can include severe agitation, confusion, hallucinations, and autonomic storm	This window requires monitored medical care — inpatient or

Timeline	What's happening?	What to Expect	What to Do
	delirum/psiko-seendem syndroom beskryf in Afdeling 1	(racing heart, high blood pressure, heavy sweating) on top of everything in the first stage	supervised detox, not home management, for anyone with a dependent use pattern.
Days 3-7	Acute danger passing if medically managed; GABA-B system still severely depleted	Continued anxiety, insomnia, and physical discomfort as the acute crisis resolves; still requires careful monitoring.	No sudden reduction of any bridge medication (baclofen or benzodiazepine) during this window — the taper from here needs to be gradual, per Section 8.
Weeks 1-4	GABA-B reseptor upregulering beginnende; taper (indien nog aan die gang is) voortgaan onder mediese toesig	Geleidelike stabilisering; angsgolwe; slaap verbeter stadig	The full supplement stack begins (Section 9). This is also when genomic testing, if pursued, becomes actionable (Section 5).
Maande 1-3	GABA-B receptor recovery is genuinely still in its early-to-middle phase — this system upregulates more slowly than GABA-A.	Anxiety waves (a documented PAWS pattern for GHB specifically) requiring active management, not just waiting out	Consistent breathwork, exercise, and the supplement protocol in Section 9 are what determine how severe these PAWS waves are.
Maande 3–12	Die sentrale GABA-herstel fase – regtig 3–6 maande, langer as die meeste ander stowwe in hierdie biblioteek.	Gradual, real improvement; PAWS waves decreasing in frequency and intensity	Keep every protocol. This is not a phase to taper off support early.
Maande 12-18+	Substantial GABA-B receptor recovery reaching completion	'n regtig ander basislyn — nie net simptoomafwesigheid nie, maar 'n senu-stelsel wat weer op sy eie regulasie loop	Dis is 'n regte transformasie, nie net tyd wat verbygaan nie. Dit vereis gebou word deur die protokolle

Timeline	What's happening?	What to Expect	What to Do
			in hierdie gids, en nie bloot gewag word vir nie.

DIE CENTURION

Notice how much longer this timeline runs than the benzodiazepine or methamphetamine guides in this series. The GHB Guide's own material states this directly: 'unlike most other substances where the central phase is 4-8 weeks, GHB's GABA-B receptor upregulation is slower and more prolonged. The GABA restoration phase genuinely takes 3-6 months.' Expecting benzo-guide or meth-guide timelines here will set you up to feel like recovery has failed when it's actually still on schedule.

Section 3 — Hoe lank voordat die brein regtig normaal is

The honest answer here is longer than for any other substance in this series so far. The GABA-B receptor recovery that reaches substantial completion around months 12-18 is not simply symptom resolution — it represents the nervous system rebuilding genuine endogenous regulation after having outsourced that regulation to a substance dosed every 2-3 hours around the clock, for however long the dependence lasted. That's worth sitting with directly: for someone dosing that frequently, GHB wasn't just a substance being used — it was functioning as the nervous system's external operating system. Rebuilding the internal one takes real time.

BEING HEARD

As jy by maand 5 of 6 is en nog angsgolwe kry wat uit die niks kom, beteken dit nie bewys dat die herstel nie werk nie — volgens die tydlyn hierbo, is jy presies waar die biologie voorspel dat jy sal wees. Die eindstreep hier word gemeet oor meer as 'n jaar, nie weke nie. Dit is nie 'n ontmoedigende getal nie. Dit is die eerlike een, en dit beteken die golwe wat jy nog ry, is verwag, nie 'n teken van mislukking nie.

Section 4 — Wat jy waarskynlik sal ervaar

Out of the acute medical emergency window in Section 2, the recovery period has a distinct texture worth naming directly.

- Anxiety that comes in waves rather than a steady background hum—this is a documented PAWS pattern specific to GHB recovery, not a sign of relapse risk escalating on its own.
- Sleep that is genuinely difficult to rebuild, especially for anyone who was dosing overnight to manage sleep directly — the nervous system has to relearn how to fall and stay asleep without a GABA-B agonist doing that work externally.

- 'n spesifieke soort desoriëntasie wat nie net emosioneel is nie – vir langterm-, hoë-frekwensie gebruikers, het die dwelm die letterlike struktuur van die dag georganiseer (die volgende dosis was nooit meer as 'n paar uur weg nie). Dit verwyder kan 'n temporele en identiteitsdesoriëntasie veroorsaak bo-op die neurochemiese terugtrekking – nie net 'wat voel ek' maar 'wat doen ek selfs met my tyd nou.'
- Autonomic sensitivity that can persist past the acute window — a racing heart or sweating in response to ordinary stress, echoing the autonomic storm of early withdrawal at a much lower intensity.

Section 5 — Genetic Cross-Reference

Gene	Wat dit regeer	GHB Recovery Relevance
GABRA2	GABA-A reseptor alfa-2 subeenheid — basis inhiberende toon	Die enigste mees belangrike gen vir GHB kwesbaarheid. Hoërisiko variant draers loop struktureel 'n laer-krag basisbrekende stelsel, en GHB aktiveer dit kragtig ondergevoede stelsel. Hierdie profiel beteken stadiger GABA-herstel, meer ernstige PAWS, en ononderhandelbare (not optional) natuurlike GABA-bou praktyk — kyk na Afdeling 9.
GABBR1 / GABBR2	GABA-B receptor subunit genes — die primêre reseptor wat GHB self op werk	The most directly relevant genes to GHB specifically, as distinct from GABRA2's relevance to benzos/alcohol. Variants here affect individual sensitivity to GHB and the severity of GABA-B receptor downregulation with chronic use. Testing availability is currently limited, but this is worth tracking as the field develops.
COMT (Val158Met)	Dopamine and stress chemical clearance	Met/Met ('Sage,' slow clearance) is overrepresented in anxiety-driven GHB use — the drug's GABA-B anxiolysis relieving a chronically overstimulated system. Val/Val ('Warrior') appears more in chemsex/stimulant-adjacent use, with GHB functioning as the counterbalance to stimulant-driven anxiety. This determines recovery direction: Sage needs calm-building; Warrior needs stimulation-management.
MTHFR	Methylation – neurotransmitter synthesis	Creates baseline neurochemical deficits GHB was partially addressing.

Gene	Wat dit regeer	GHB Recovery Relevance
		5-MTHF supplementation is foundational and often produces early, visible improvement. GHB also depletes B-vitamins through its own metabolic pathways, making methylation support urgent rather than optional in early recovery.
DRD2 / ANKK1 (Taq1A)	Dopamine DRD2 receptor density — Reward Deficiency Syndrome	Relevant particularly in chemsex and nightlife contexts, where GHB's dopaminergic reward partially fills the RDS gap. Recovery must build dopamine reward architecture directly, or the biologically-driven pull toward resuming GHB (or any dopaminergic substance) remains.
SLC6A4 (5-HTTLPR)	Serotonin transporter expression	Short/short carriers run high anxiety sensitivity, predisposing toward GABAergic substance use for anxiety management. Recovery needs to address the serotonin system alongside GABA — 5-HTP and 5-MTHF support both.
CYP2C19 / CYP3A4	The metabolism of baclofen and benzodiazepines used in GHB detox	Direkty relevant vir Mediese Bestuur in Afdeling 8 — swak metaboliseerders akkumuleer baclofen by standaarddosisse; ultra-vinnige metaboliseerders mag hoër dosisse benodig. Informeer die behandelende arts van enige bekende status.
ADH1B / ALDH2	Alcohol and 1,4-BD metabolism	Specifically relevant for 1,4-BD users — the ADH/ALDH pathway is what converts 1,4-BD into GHB in the liver. ALDH2*2 carriers who used 1,4-BD alongside alcohol face compounding conversion-pathway conflict and acetaldehyde toxicity accumulation; liver function testing is a priority for this specific subgroup.

Who's Actually Most at Risk — Putting the Genes Together

Two separate questions get asked about genetics here, and they have different answers worth keeping distinct: who is most likely to develop dependence in the first place, and, separately, whose withdrawal and taper will be hardest once dependence exists.

Question	Die Gene wat dit werklik beantwoord	Hoekom
People who have a history of substance abuse, those with existing mental health conditions, and individuals who use GHB frequently are most at risk.	GABRA2 high-risk + COMT Met/Met ('Sage')	This is the same 'Hard Fail' combination the compendium identifies for GABAergic substances broadly: a structurally underpowered baseline braking system (GABRA2) combined with a chronically overstimulated, hard-to-quiet nervous system (Met/Met). GHB's rapid, complete anxiolysis is disproportionately compelling for exactly this profile — it's not that this person has less willpower, it's that GHB is answering a louder, more specific biological question for them than it is for someone without this combination.
Whose withdrawal and taper will be hardest, regardless of why they started?	GABRA2 high-risk + GABBR1/GABBR2 variants	These are the receptor-level genes, not the motivation-level ones. GABRA2 determines general inhibitory reserve; GABBR1/GABBR2 determine sensitivity and downregulation severity at the exact receptor GHB acts on. This profile predicts a genuinely slower taper and more severe PAWS almost independent of the person's psychological relationship to the drug.

⚠ WAARSKEUWING

A direct answer to the COMT/taper question specifically: COMT genotype does not appear to determine whether a taper mechanically works — that's governed by the GABA receptor genes above, not by dopamine clearance speed. What COMT actually determines is why someone used and what direction their post-taper recovery needs to point. A Met/Met ('Sage') profile tapering off GHB needs the taper itself managed the same way as anyone else with their GABRA2/GABBR status — but their longer-term recovery work is calm-building (the anxiety was there before GHB and needs a real answer). A Val/Val ('Warrior') profile, more often found in chemsex/stimulant-

counterbalance use, needs stimulation-management instead — genuine high-intensity outlets, not just calm. Same taper mechanics either way; different recovery target afterward.

Section 6 — Special Populations

Situation	What Makes It Different
Sodium oxybate prescribed for narcolepsy with cataplexy	Sodium oxybate (pharmaceutical GHB) is the standard of care for this condition, and the dependence that develops is a necessary consequence of appropriate treatment, not a moral failure or a sign of misuse. Discontinuing it requires neurologist involvement, because the narcolepsy symptoms being managed will return on cessation. Alternative narcolepsy treatments (modafinil, armodafinil, solriamfetol, pitolisant) should be optimised before or alongside any reduction. This is a clinical decision for a specialist, never a self-managed taper.
GHB used as an intentional alcohol substitute	Sommige menne gebruik GHB spesifiek om te drink te verminder – hulle ruil een GABAerge substansie vir 'n ander met 'n ander risiko-profiel. GHB mag laer orgaan-toksisiteit dra as swaar alkoholgebruik by gematigde dosisse, maar sy afhanklikheidsverantwoordelikheid en oordoseringrisiko is op hulle eie ernstig. Indien dit u situasie beskryf, kan beide afhanklikhede saam bestaan, en 'n mediese assessering van gekombineerde GABAerge afhanklikheid is noodsaaklik voordat enige vermindering van een of twee substansies poging word.
Growth hormone (GH) supplementation for bodybuilding and athletic enhancement remains a highly controversial topic, with significant scientific debate surrounding its efficacy, safety, and appropriate usage. **Mechanism of Action:** GH plays a crucial role in normal human development, metabolism, and tissue repair. It stimulates the liver to produce Insulin-like Growth Factors (IGF-1), which are	GHB stimule die pituitary groeihormoonvrystelling by lae, nie-rekreasionele dosisse, en word deur sommige bodybuilders en atlete spesifiek vir hierdie effek gebruik — dikwels elke nag geneem, ooreengekom as 'n herstel- en spierbouhulp eerder as as 'n partydrug. Dit is 'n algemene en ondergewerkte beginpunt vir GHB-afhanklikheid presies omdat dit nie lyk soos rekreasionele dwelmgebruik vir die persoon wat dit doen nie: dit word alleen geneem, op 'n rooster, vir 'n gespesifiseerde prestasiedoelwit, wat kan vertraag om die patroon as afhanklikheid vorm te herken totdat slaap chemies afhanklik daarvan geraak het. Die nagtelike doseringstruktuur self is dieselfde meganisme described in Section 1 and Section 4 — a habitual, scheduled dose is exactly how tolerance and physical dependence establish, regardless of the stated reason for taking it. Recovery needs to address the growth hormone axis directly rather than just the GABA system — natural GH-stimulating strategies (deep sleep architecture, appropriately timed fasting, HIIT training) replace the pharmacological approach; physician-guided GH-secretagogue peptides (such as ipamorelin) have been used as a bridge in some cases. This population also frequently needs a specific reframe: a supplement taken for a legitimate athletic goal becoming a dependence is not a different

Situation	What Makes It Different
responsible for many of GH's anabolic effects. In theory, supplementing with exogenous GH could boost muscle protein synthesis, improve recovery rates, and enhance overall body composition. **Potential Benefits Claimed by Users:** * Increased lean muscle mass accretion. * Improved fat loss (lipolytic effect). * Enhanced athletic performance and recovery. **Scientific Consensus and Concerns:** While anecdotal evidence is plentiful in bodybuilding circles, the scientific literature presents a more nuanced picture: 1. **Natural Endogenous Production:** The body's natural pulsatile release of GH is tightly regulated by the pituitary gland. Exogenously administering high doses can disrupt this delicate feedback loop. 2. **Side Effects:** Over-stimulation carries risks, including: * Acromegaly (enlargement of hands,	or lesser kind of addiction than any other in this guide — the biology doesn't check the stated intent.

Situation	What Makes It Different
feet, and facial features). * Insulin resistance and potential for developing Type 2 Diabetes. * Joint stress and ligament damage due to rapid tissue changes. **Usage in Athletics:** In competitive sports, the use of GH is generally classified as a performance-enhancing drug (PED) and is banned by most anti-doping agencies (e.g., WADA). Athletes seeking enhancement must weigh potential gains against severe health risks and career repercussions. ***Disclaimer: This information is for educational purposes only and does not constitute medical advice. The use of GH or any similar substance should only be undertaken under the direct supervision of a qualified endocrinologist or sports physician.***	
Dance festivals and rave culture	Historically one of the most common entry points to GHB use of all — a euphoric, sociable dose shared at a party or festival, often alongside MDMA, ketamine, or alcohol, used to enhance the experience or ease the comedown from stimulants. This is also the context where Section 1's 'unpredictability' danger is most acute in practice: festival and club GHB is typically measured in capfuls poured from an unlabeled bottle, shared peer-to-peer with no way to know the actual concentration, on top of a person's own

Situation	What Makes It Different
	uncertainty about how much they've already had that night. Concentration inconsistency between one bottle and the next is not a rare edge case in this context — it is close to the norm, which is a major reason GHB overdose events cluster heavily around festival and club settings specifically. The transition into dependence here often isn't a single dramatic moment; it's using between events to chase the same social ease or sleep support the festival dose provided, with frequency quietly climbing as festival season turns into a regular circuit. Alcohol co-use is also especially common in this specific context, given the setting — see Section 5's 'Hard Fail' combination note.
Livestreaming en sosiale media-inhoudskepping	A genuinely newer context than the others in this table, worth naming on its own rather than folding into 'nightlife' generally. Using on camera, for an audience, adds a second reward system stacked directly on top of the pharmacological one — real-time viewer attention and engagement activate the same dopaminergic reward circuitry the substance itself is activating, and the two can reinforce each other in a way that plausibly accelerates escalation compared to private use. This context also inverts the safety logic the rest of this guide depends on: a bystander who is physically present can recognise an overdose and act — help skree, plaas iemand in die herstelposisie, verwyder ander stowwe van bereik. 'n Gehoor wat na 'n livestream kyk kan niks hiervan doen nie. Sigbaarheid is nie dieselfde as veiligheid nie, en 'n kamer vol kykers is, funksioneel, dieselfde as om alleen te wees. Moderne GHB of GBL word ook toenemend in vermomde houers gelewer — energie-drankbottels, sportdrankbottels — soms deur 'n derde party eerder as gemeter deur die persoon wat dit gebruik, wat Seksie 1 se konsentrasie-onvoorspelbaarheidsgevaar met 'n tweede laag vererger: nie weet wat eintlik in die bottel is nie, boopop. of not knowing its strength.

BEING HEARD

There's a pattern worth naming that doesn't fit neatly into the table above because it isn't really about why someone uses GHB — it's about where. When a venue, a club, a friend group, or a community bans or excludes someone because of their use, the use itself very rarely stops. What changes is where it happens. A person who was using around other people — however imperfectly supervised, however far from ideal — often shifts to using alone, at home, specifically because the social settings that once absorbed the behaviour no longer will. That shift from 'someone nearby who 'No one around at all' is nie 'n klein detail nie. Dit is die presiese veranderlike wat bepaal of 'n oordos waargeneem en oorleefbaar is of stilweg gebeur. Uitsluiting, ongeag sy redes, ruil dikwels 'n sigbare probleem vir een onsigbaarder en gevaarliker. As dit jou eie geskiedenis beskryf — 'n verbanning, 'n rissiwaring, 'n gemeenskap wat 'n deur gesluit het — dan was daardie isolasie nooit 'n teken dat jy dit alleen moes hanteer nie. Dit was

'n strukturele gevolg van waar die ondersteuning gewyk het, nie 'n vonnis oor of jy enige verdien het nie.

Section 7 — Use Pattern Classification

Die meeste van hierdie gids, uit noodsaaklikheid, spreek die gevaarlikste kant van GHB-gebruik aan – want daardie kant kan doodmaak, en dit deel regkry is belangriker as enigiets anders hier. Maar die meeste mense wat GHB gebruik, is nie by daardie punt nie, en iemand wat een keer per week gebruik en bloot wil stop, het 'n werklik ander antwoord nodig as om na 'n noodafdeling te gaan,' nie net 'n dunner weergawe van dieselfde waarskuwing nie.

Level	Pattern	Risk of Dependence/Withdrawal	What Stopping Actually Requires
Rare / experimental	A handful of times total, no regular pattern	Very low	Fysieke afhanklikheid is eintlik nie 'n kommer op hierdie vlak nie. As jy wil ophou, kan jy net stop. Die belangrikste ding om te doen is om te verstaan waarom dit jou aangetrek het (kontekstetaakel van Afdeling 6) sodat 'n vervangende trek nie stilweg ergens anders vorm nie.
Occasional / social	Roughe monthly or less, tied to specific contexts (parties, chemsex, etc.), non-escalating over time	Laag	Stopping alone is reasonable for most people at this level. Worth watching for: has the interval been shrinking? Has the amount per session been creeping up? If either is true, you may be sliding toward Level 3 rather than staying at Level 2.
3 — Regular recreational, not dependent	Roughly weekly, stable rather than escalating, no nocturnal dosing, no physical withdrawal symptoms noticed between uses (no shakiness, anxiety spike, or sweating before the next planned use)	Moderate but generally manageable without medical detox	This is the population most missing from most GHB material, including earlier drafts of this guide. Stopping is very likely safe to do without inpatient detox — but 'likely safe' isn't 'certain,' so a same-day

Level	Pattern	Risk of Dependence/Withdrawal	What Stopping Actually Requires
			conversation with a physician before stopping is still the responsible move, particularly to rule out dependence you may not have noticed forming. Section 9's Tier 1 and Tier 3 (nutrition and natural GABA support) are directly usable here, starting immediately rather than after a medical detox phase.
4 — Eskalerend / afhanklikheidsvorming	The frequency or dose has been increasing; any physical discomfort noticed between doses (anxiety, shakiness, disrupted sleep) even if not yet daily.	Hoog en aktief stygend	Dit is die vlak waar mediese ingryping voordat stop nie meer opsioneel voorsigsaamheid word nie en 'n werklike veiligheidsvereiste word — selfs sonder daaglikse gebruik, eskalasie self is die waarskuwingsteken dat die GABA-B stelsel aanpas. Wag nie vir Vlak 5 om mediese leiding te soek nie.
5 — Afhanklik (die nooddrempel)	Drie of meer kere per dag, insluitend nagtoedings om slaap te bestuur	Severe — this is explicitly the threshold Section 0 identifies as a medical emergency requiring supervised detox	Do not attempt to stop without medical supervision under any circumstance. This is the population Section 8's detox protocol is written for.

SANCTUARY

As you're at Level 2 or 3 and reading this whole guide's emergency language wondering if it applies to you — it's worth naming directly: it mostly doesn't, and that's a good thing, not a reason to dismiss the guide. The seriousness in Section 0 exists for Level 5, and it needed to be said clearly and first because that's where the mortality risk actually lives. But wanting to stop a weekly, non-dependent pattern is a completely legitimate reason to be here, and it doesn't require the machinery of Section 8 to do safely —

mostly it requires the same physician check-in any responsible change deserves, plus the natural rebuilding work in Section 9, which works just as well starting from a lighter baseline as a heavy one.

Hoarding a Supply: A Risk Factor Independent of Use Level

This applies across every level in the table above, not just the dependent end: keeping a large stored supply on hand is its own distinct risk factor, separate from how often someone is actually using. Simple availability changes behaviour — a large supply removes a natural point of friction that a smaller, harder-to-replace amount provides. In an already-intoxicated state, with judgment already affected, having more within reach makes an impulsive extra dose far easier than it would be if getting more required leaving, waiting, or making contact with someone. This is the same mechanism environmental redesign addresses elsewhere in the Recovery Kernel (Compendium Protocol 05) — access itself is part of the risk architecture, not just the substance.

⚠ WAARSKEUWING

The danger compounds specifically with GHB because of what Section 1 already establishes about concentration inconsistency. A large hoarded supply — especially GBL or 1,4-BD, or GHB from an unlabeled source — carries no guarantee that concentration stays consistent from the first measure to the last. Redosing repeatedly from the same large stockpile, already intoxicated and already less able to judge cumulative dose accurately, is a documented pattern behind escalation and overdose alike. A large supply on hand doesn't just make more use possible — it makes the specific kind of dosing error this entire guide exists to warn about more likely.

As anything, reducing or stopping is the goal at any level in the table above, reducing the accessible supply itself — not just the intention to use less — is a concrete, practical step worth taking on its own, separate from anything pharmacological.

Section 8 — Die Mediese Ontgiftingsprotokol

Hierdie afdeling bestaan om jou of iemand wat jou ondersteun te help om vir die regte sorg aan te spreek – GHB-onttrekkingbestuur word werklik onderbedien in spoedgeneeskunde, en vooraf weet van die opsies kan die verskil maak in wat 'n behandelende span bied.

Approach	Mechanism	Clinical Notes
Baclofen taper (preferred where available)	GABA-B agonist — dit vervang direk vir GHB op die reseptorvlak; langer halflengte stel geleidelike, stabiele afbou toe	Die mees farmakologies raasionale benadering. Die vereiste dosisse is dikwels baie hoog — tot 200-300mg/dag — en moet noukeurig getitreer word. Nie alle nooddepartemente is bekend met baclofen vir hierdie spesifieke gebruik nie; jy mag dit direk moet

Approach	Mechanism	Clinical Notes
		bepleit, deur die kliniese literatuur aan te haal (Addolorato et al. 2009; Bell et al. 2011) indien daar onbekendheid mee gelei word.
High-dose benzodiazepine protocol (most commonly used in practice)	GABA-A agonist — addresses die GABA stelsel breedvoerig, by 'n ander reseptorondersteuntyf as GHB se primêre teiken.	Often used simply because it's familiar to emergency physicians. Less pharmacologically specific than baclofen and may require very high doses to control severe withdrawal, with a real risk of transitioning into benzodiazepine dependence on top of the GHB Recovery. Requires close medical monitoring.
Sodium Oxybate tapering	Direct substitution with the same molecule in controlled, measured form, then gradual taper	Die mees teoreties elegante benadering – dieselfde molekuul, bekende farmakokinetika, geen raaiery oor voorsieningskonsentrasie nie. Word in sommige spesialist-verslawingseenhede gebruik; beskikbaarheid is beperk buite spesifieke kliniese kontekste en vereis 'n spesialiste verslawingsarts.
Phenobarbital	Langwerkende barbituraat met GABA-A agoniste-aktiwiteit; ekstreme halwe lewensduur (53-118 ure) bied inherent stabiele, self-afnemende plasmaniveaus	Gebruik in gespesialiseerde omstandighede vir komplekse GABAerge polidrug-onttrekkingssyndroom. Vereis noukeurige respiratoriese monitering. Nie eerste-lyn nie, maar waardevol in refrakterie gevalle.
Supportive medications (adjunctive to any of the above)	Clonidine vir die outonomie storm; antipsikotika (haloperidol, quetiapine) vir delirium/hallingenasie; antikonvulsante vir krampsvoorkoming; IV vloeistof en elektrolytebestuur	GHB withdrawal produseer ekstreem outonomiese instabiliteit saam met psigiatriese simptome — hierdie bykomende behandelings spreek direk die spesifieke simptoomklusters in Afdeling 1 en Afdeling 2 aan.

⚠ WAARSKEUWING

A note on active research: there is real, published toxicology research into monocarboxylate transporters (MCTs), which mediate GHB's renal reabsorption, and into whether urinary alkalinization (pH manipulation, including via sodium bicarbonate)

can modestly enhance GHB renal elimination — the same underlying principle used clinically for salicylate overdose. A GABA-B antagonist (SCH-50911) has shown reversal effects in animal studies but has not been tested in humans. Both are described in the human proof-of-concept study cited in the literature as exactly that — proof of concept, not clinical practice. This is a genuine and active research frontier, worth knowing exists. It is not a home intervention, not a substitute for the protocols above, and this guide will not walk through dosing or administration for either — that specific gap between 'the mechanism is real' and 'this is safe to do yourself, unmonitored' is exactly where people get hurt.

When Hospital Access Is Genuinely Limited

Not everyone reading this can reach a hospital — because of distance, cost, immigration status, a country with limited emergency infrastructure, or a dozen other real reasons. That gap deserves a real answer, and the honest one isn't a self-administered protocol — it's making sure the resources that exist specifically for this situation actually get used.

What's Actually Available Without a Hospital	Hoekom Dit Saakmaak
Poison Control hotlines	Die meeste lande het 'n gifbeheer- of toksikologiehotline wat per telefoon bereikbaar is en deur klinici bemanning word wat spesifiek opgelei is om presies hierdie soort situasie op afstand te trias – insluitend gevalle waar vervoer nie onmiddellik moontlik is nie. Dit is real-time kundige leiding, nie 'n dokument nie. Om een te bel is 'n betekenisvol anders en veiliger optrede as om slegs skriftelike materiaal te volg, selfs van afstand en selfs al kan 'n hospitaal werklik nie vinnig bereik word nie.
Tele-health, waar beskikbaar	'n real-tyd video of telefoonverbinding met 'n dokter, selfs een wat nie fisies teenwoordig is nie, stel werklike kliniese oordeel toe — die assessering van bewustheid, asemhaling en reaksie op simptome soos dit verander — op 'n manier wat geen statiese protokoll kan nie.
Never alone during withdrawal from a dependent-level use pattern	As long as a hospital is genuinely out of reach, the single highest-value thing available is a sober person physically present who can monitor breathing and consciousness, place someone in the recovery position if needed, and call for whatever help exists the moment something changes — not to administer treatment, but to be the early warning system a monitor would be in a hospital.
Acting before the emergency threshold, not during it	As limited access to care is a known reality — not a hypothetical — the highest-leverage moment is before use reaches the dependent, three-or-more-times-daily pattern in Section 7, when a supervised reduction is still possible through whatever care access does exist, rather than after withdrawal has already begun without

What's Actually Available Without a Hospital	Hoekom Dit Saakmaak
	anywhere to go.

BEING HEARD

As jy hierdie lees omdat 'n hospitaal regtig nie 'n realistiese opsie vir jou is nie, wil ek direk sê: dit is 'n mislukking van die stelsels en omstandighede rondom jou, nie iets wat hierdie gids kan oplos deur jou 'n protokolo te gee nie. Wat dit wel kan doen, is om jou na die telefoongebaseerde hulpbronne te wys wat bestaan vir presies hierdie gaping, en na vroeër optree, terwyl meer deure nog oop is, eerder as later wanneer minder oop is. Dit is 'n regte antwoord, selfs al is dit nie die een wat jy gehoop het nie.

Die 30-Dae Afgrond: 'n Tweede Hervalingsrisiko-As Ontslagbeplanning Wat Dikwels Misgaan

⚠ WAARSKEUWING

A common discharge pattern deserves to be named directly, because it creates a second, distinct danger window that isn't the same as the acute withdrawal covered above: inpatient management for GHB withdrawal (often around 5 days) followed by discharge with a benzodiazepine supply — commonly around 30 days — intended as a bridge while the GABA-B system recovers. When that 30-day supply runs out, there is a documented, high incidence of relapse at exactly that point. This is not a failure of willpower or of the discharge plan being followed. It's a foreseeable pharmacological event.

Die rede pas direk by Seksyen 3 se tydlyn: op dag 30 is GABA-B reseptorherstel nog in sy vroeë fase – werklik maande weg van die substansiële herstel wat hierdie gids se eie data op 12–18 maande plaas. Die benzodiazepinebrug het nooit net simptome gedurende daardie venster behandel nie; dit het GABA-A ondersteuning gebied wat gedeeltelik kompenseer vir 'n GABA-B stelsel wat nog nie herbou is nie. Wanneer die voorskrif op 'n vaste kalenderdatum en nie op 'n kliniese een nie eindig – dikwels sonder afbou, en dikwels sonder enige ambulante opvolging geskeduleer voordat die voorraad op is – ervaar die persoon 'n tweede onttrekkingssyndroom. landing directly on top of a first one that was never actually finished. The chronic rebound anxiety and fear this produces is neurobiological, not primarily psychological, which is exactly why supportive therapy alone is frequently insufficient here: talk therapy is the right tool for a different problem than an unrecovered GABA-B receptor system generating genuine, physiologically-driven anxiety.

What Discharge Planning Often Gets Wrong	What Actually Reduces This Risk
Treating die 30-day supply as the end of the recovery timeline rather than a small piece of a 12-18 month one	Scheduling an outpatient follow-up before discharge, with a specific appointment date well before the medication runs out—not 'follow up with your doctor,' but a booked date.

What Discharge Planning Often Gets Wrong	What Actually Reduces This Risk
A hard stop on the benzodiazepine at day 30 rather than a taper	'n Ware afbou van die brugmedikasie, gekoördineer met 'n verslawingsgeneeskunde of psigiatriese voorsitter, wat dieselfde persentasiegebaseerde verminderingsbeginsel volg wat deurgaans in hierdie biblioteek gebruik word eerder as 'n vaste-hoeveelheid voorskrif wat net ophou.
Assuming supportive therapy alone is sufficient for the anxiety that follows	Recognising that rebound anxiety at this stage has a real physiological driver (Section 3) and may require the Tier 1 and Tier 3 support in Section 9 specifically, not just talk therapy, alongside any needed psychiatric medication management.
No explicit warning to the patient that day 30 is a known high-risk point.	Naming this risk directly, before it arrives — which is the entire purpose of this subsection existing. Knowing a cliff is coming changes how a person approaches it.

SANCTUARY

As you're nearing the end of a discharge prescription right now, or if you've relapsed after one ran out, listen to this plainly: what happened at that point was not a personal failure of willpower, and it doesn't erase any recovery work done during the inpatient stay or the 30 days that followed. It was a foreseeable gap in how the medication timeline and the biological timeline were matched up. The right response is the same as anywhere else in this guide facing a real risk — get back to medical support and a real taper plan, not a verdict. whether recovery is possible for you.

How This Can Become a Second, Separate Addiction

There's a specific pattern worth naming plainly, because it's less about the substance and more about what happens when nobody explains what's actually happening inside a person's own nervous system. Someone whose discharge benzodiazepine has run out is often still experiencing real, physiologically-driven anxiety — the direct product of a GABA-B system that, per Section 3, is nowhere near recovered. If nobody has told them this is what day 30 feels like and why, the natural conclusion is not 'this is expected post-GHB recovery.' It's 'I have an anxiety disorder' — a completely reasonable misread of a real symptom, made without die een stukkie inligting wat alles sou verander oor hoe dit behandel moet word.

What often follows is a visit to a psychiatrist, sometimes with the GHB history disclosed and sometimes not — disclosure can feel harder than the symptom itself, given the stigma this library has already named elsewhere. A psychiatrist evaluating what presents as straightforward, severe anxiety, without the full context of a recent GABA-B withdrawal in progress, is often making a reasonable clinical call by prescribing a benzodiazepine. It isn't malpractice, and the person

isn't necessarily lying — they're describing something genuinely severe and getting a response that matches what they described, minus the context that would have changed the response. This is the same pattern the Recovery Kernel names directly elsewhere as 'withdrawal framed as relapse': a temporary, biologically-driven post-acute symptom gets treated as a standalone, chronic condition, and the medication prescribed for the chronic version has no natural end date — because nothing in the visit suggested one was needed.

⚠ WAARSKEUWING

Die resultaat is 'n nuwe, dikwels langer-voortdurende afhanklikheid as die oorspronklike een: 'n benzodiazepine wat onbeperkt voorgskryf word vir wat eintlik 'n tydsbeperkte venster van post-GHB neurologiese herstel was. Dit is nie 'n hipotetiese risiko nie — dit is 'n gedokumenteerde patroon, en dit verdien dieselfde gewig as die GHB-afhanklikheid wat hierdie hele gids aanspreek, want farmakologies is dit dieselfde val in 'n ander bottel.

What actually reduces this risk: disclosing the recent GHB history and withdrawal timeline to any psychiatrist or prescriber involved, even though it's uncomfortable, specifically because it changes what an informed prescriber would recommend. A prescriber who knows they're looking at post-GHB GABA-B recovery, not a new primary anxiety disorder, can reasonably offer a short-term, clearly time-limited bridge with a stated review date — the same accountability standard named elsewhere in this library for any substitute medication — rather than an open-ended prescription. The exact language worth using: 'This anxiety started after I stopped GHB about a month ago, and I Understand my nervous system might still be recovering from that specifically—I'd like whatever we try to have a clear review date rather than being open-ended.' That sentence alone can be the difference between a real bridge and a second lifelong dependence.

Section 9 — Die Herstelprotokol (Post-Akut)

⚠ WAARSKEUWING

This section starts once the acute medical emergency window (Section 0, Section 2's Hours 12-72) has passed and a doctor has confirmed you are medically stable. It doesn't replace Section 8—it's what follows.

Tier 1 — Biologiese Grondslag (Dag 1 na ontgiftingsproses voort)

Support	Role
Magnesium glycinate (400-600mg, split doses)	GABA-A reseptor ondersteuning en algemene senuwstelsel kalmering – fundamenteel, nie opsioneel nie, vanaf Dag 1.
P5P (active B6, 25-50mg)	Cofactor vir die glutamaat-na-GABA-omskoming (glutamaatdekarboksilase) — direk relevant gegewe die glutamaatstoot in vroeë terugtrekking (Afdeling 2).
5-MTHF (400mcg)	Foundational methylation support, especially crucial

Support	Role
	given GHB's own B-vitamin depletion through its metabolic pathways.
L-theanine (200-400mg as needed)	Non-sedating GABA/glutamate modulation vir anxious waves without dependence liability.

Tier 2 — Mediese en Psigiatrië Ondersteuning

- Buspirone (non-addictive serotonin 1A partial agonist) vir aanhoudende post-detoks angs – neem 2-4 weke vir volle effek, geskik vir die volgehoude angs van hierdie herstel sonder die risiko van benzodiazepine-afhanklikheid.
- 5-MTHF trial before SSRI consideration for MTHFR-variant depression – assess response over 4–8 weeks; SSRIs remain appropriate if clinically indicated.
- Low-dose quetiapine (25-50mg, time-limited) kan slaaparsitektuur ondersteun by ernstige GHB-verwante slapeloosheid wanneer die aanvullingsstap alleen onvoldoende is — nie eerste-lyn nie, maar 'n regte opsie.

Tier 3 — Nervous System Regulation (Weeks 1-12, die sentrale herstel fase)

- Breathing exercises—daily, non-negotiable given the extended GABA-B recovery timeline in Section 3.
- Kou blootstelling en yoga – albei direk relevant vir die opbou van natuurlike GABA-toon.
- Consistency is more important here than intensity — according to Section 2's timeline, this is a 3–6 month central phase, not a 4–8 week one. What determines PAWS severity is consistent practice throughout that whole period, not a strong start for the first two weeks.

Exercise Specifics — Hoekom Kardio Hier Meer Saak Is As In Die Meeste Ander Gidsies

GHB withdrawal behels regte autonome instabiliteit (Afdeling 1, Afdeling 2) — hartsklop wat te vinnig klop, bloeddrukswings, swaar sweet – en kardiovaskulêre oefening is direk relevant vir die herstel van die onderliggende stelsel, nie net 'n algemene welstandsaanbeveling nie.

Type	Wat dit produseer	Frequency
Aerobics (hardjogging, bike riding, swimming, fast walking)	Endorphins, serotonin, BDNF — en spesifiek hier relevant, die geleidelike normalisering van rus outonome toon (hartfrekwensievariabiliteit, basis bloeddruk) terwyl die kardiovaskulêre stelsel herstel van die onttrekkings-era outonome storm.	3-5 sessions per week
Strength training	Dopamine, testosterone, growth hormone, BDNF — relevant particularly for the	2–4 sessions per week

Type	Wat dit produseer	Frequency
	athletic/bodybuilding use context in Section 6, where GHB's growth-hormone-stimulating effect was part of the original appeal and needs a genuine replacement	
Yoga	Parasympatiese aktivering en direkte angsvermindering – 'n GABA-ondersteunende praktyk op sy eie, nie net 'n aanvulling by asemhalingsoefeninge nie.	1–3 sessions per week
Daglike stap	Die nie-onderhandelbare basislyn – lae drempel, werklik effektief, gepas selfs op die vroegste en mees uitgeputte dae van herstel.	Daily

⚠ WAARSKEUWING

Timing en intensiteit is belangrik, veral aan die begin. Oortraining in die eerste weke voeg fysiologiese stres by op 'n reeds belas stelsel eerder as om dit te verlig — begin met stap en sagte beweging, bou na die volledige program in maande 1-3, en hou intensiewe oefening spesifiek nie in die aand nie, aangesien dit die slaapbegin vertraag in 'n herstel wat reeds vir slaapgehalte veg (Afdeling 4).

Nutrition Scaffolding — Beyond the Core Four Supplements

Tier 1 se vier supplemente is fundamenteel, nie die hele prentjie nie. GHB-gebruik – veral swaar, kroniese gebruik – ontwrig voedingsstatus wyd, en die senuweestelsel wat die herstelwerk beskryf word deur hierdie gids benodig, het die rauwe materiale nodig om dit mee te doen.

- Consistent protein intake supports neurotransmitter synthesis broadly, including the GABA and dopamine systems this entire recovery depends on.
- Consistent mealtimes are just as important as what you eat during early recovery—blood sugar dips worsen the anxiety and mood swings that are already there.
- Hydration en elektrolyte ondersteuning spreek direk die outonome/swet-symptome in Afdeling 2 en Afdeling 4 aan.
- B-vitamine repletion broadly (not just P5P and 5-MTHF) is relevant given GHB's own metabolic depletion of this vitamin family — a general high-quality B-complex alongside the targeted Tier 1 items is a reasonable addition.

Losing Interest in the Drug: Natural GABA Reconstruction

Everything above manages symptoms and rebuilds biology. This part is about something different — actually wanting it less, not just tolerating its absence. GHB gave a specific thing: not partial relief, but a nervous system that would finally, completely stop. The premise here is that this exact state is genuinely reachable without it, through the brain's own systems, and that a reconstructed version is more sustainable than the pharmacological one ever was — because it doesn't downregulate the way GHB's receptor did.

Natural Activation Method	How Strong the GABA Effect Is	GHB-Recovery-Specific Notes
Uitgebreide uitademing asemwerk (4-7-8, blokasemhaling, koherente asemhaling)	Hoog – direkte vagale/parasimpatiese /GABAerge aktivering	Die enigste mees belangrike natuurlike GABA-instrument in hierdie hele gids. Dit is nie 'n metafoor nie — verlengde-uitasem asemhaling verhoog meetbaar die GABA-toon en verminder kortisol binne minute. Dae-naal praktyk totdat dit net so outomaties is soos die ou doseringsrooster was.
Koue water blootstelling	Hoog – parasimpatiese aktivering plus 'n daaropvolgende volgehoue kalmte na die aanvanklike skokreaksie.	A cold shower, 30-90 seconds, building gradually to full cold. A genuine, physically real analogue to the 'system finally stops' state GHB provided.
Strenuous cardio exercise	Moderate-high, kumulatief met konsekwentheid	Kyk na die kardio-taak wat hierbo staan – dit is dieselfde oefening wat dubbele plig verrig vir kardiovaskulêre herstel en natuurlike GABA/endorfien-aktivering.
Music at 60-70 BPM	Moderate – measurable parasympathetic entrainment effect	Handig spesifiek tydens hunkering en voor slaap; lae drempel en onmiddellik toeganklik.
'n veele eisevereende kreatiewe of vloei-toestand praktyk	Moderate-high tot full attentional engagement rather than chemical suppression	Totale absorpsie tydens 'n uitdagende aktiwiteit is, meganisties gesproke, 'n natuurlike GABA-toestand – die angstige gemoed kalmeer deur betrokkenheid eerder as deur mag.

BEING HEARD

This is not about willpower, and it's not some consolation prize for not having the drug anymore. What GHB was doing from the outside, a recovered system can genuinely do from the inside — that's not a hopeful reframe, it's the actual mechanism this whole guide has been describing. The state you're chasing was always a real, accessible nervous-system state. GHB was just one way to reach it — a fast, dangerous, downregulating way. The methods in this table reach the same place through a system that gets stronger with use instead of weaker.

Section 10 — Emergency Recognition, Restated

MEDICAL EMERGENCY

GO TO AN EMERGENCY DEPARTMENT IMMEDIATELY FOR: enige tekonstiwiteit met aanvalaktiwiteite — verwarring, desoriëntering of halusinasies — extreme agitering met 'n hartsklop wat te vinnig is, hoë bloeddruk, of swaar sweet — verswakking van trilling.

Dit geld gedurende die eerste 72 uur van ophou vir enige afhanklike gebruiker, en op enige stadium daarna indien hierdie tekens optree. Vir angsgolwe, slaapmoeilikheid, en die desoriëntering beskryf in Afdeling 4 sonder enige van die bogenoemde — hierdie word verwag volgens die tydlyn, en kontak met 'n dokter dieselfde dag is gepas maar dit is nie op sigself 'n ER-vlak noodsituasie nie.

Section 11 — Full Protocol Cross-Reference Map

Phase	Protocols	Focus
Medical Safety (Days 0-7 for dependent users)	Emergency department / inpatient detox — outside the Compendium's scope	Seizure prevention, baclofen or benzodiazepine protocol, delirium management, vital sign monitoring. This guide's Section 9 and beyond begin only once this phase is medically complete.
Biological Foundation (Days 1-14 post-detox)	Compendium Protocols 07, 09, 17, 18	Magnesium, B6, methylation stack, slaaparsitektuur, voeding, darmherstel — die bemiddelende toestande vir GABA-B stelselherstel, nie opsionele bykomste nie.
Genomic Mapping (Weeks 1-4)	Compendium Section 15, Protocol 11	GABRA2 assessment specifically — die enigste mees belangrike genomiese bevinding vir hierdie herstel, wat afhanklikheidsernst, onttrekkingsernst en PAWS-ernst verduidelik.
GABA Restoration (Weeks 1-12, die sentrale fase)	Compendium Protocol 04, Protocol 01 craving interrupt	Breathwork, cold exposure, yoga, magnesium stack, anxiety management, sleep architecture — genuinely 3-6 months, not the 4-8 week timeline of most other substances in this library.
Behavioural and Environmental Architecture (Weeks 2-12)	Compendium Protocol 02, 05, 08, 13	Cue removal, addiction transfer prevention, social architecture redesign — chemsex-konteks omgewingsgebeure (plekke, apps, sosiale netwerke) is onder die kragtigste aanwysing-geaktiveerde hunkering-omgewings wat gedokumenteer is; digitale omgewingsherontwerp is net so belangrik as fisiese.
Chemsex Recovery, if applicable (Maande 1-12)	Compendium Protocol 35S series, sexual health protocols	HIV/STI testing and prevention, sexual health in recovery, sober LGBTQ+ community building — see Section 6 above.
Identity and Meaning (Maande 2-12)	Compendium Protocol 20, 21, 23, C-28	Who you are when a dosing schedule every 2-3 hours is no longer the organising structure of your day —

Phase	Protocols	Focus
		directly addressing the temporal/identity disorientation named in Section 4.
Die Lang Termyn (Maande 6-18+)	Compendium Protocol 24, 25, C-28, C-31	Purpose architecture and a fully operational natural GABA system — per Section 3, a genuine transformation that has to be built towards, not simply waited for.

Section 12 — Closing

SANCTUARY

This is die hardest withdrawal in this entire series, and the recovery timeline is the longest — over a year to substantial completion, not weeks. That's not meant to discourage. It's meant to make sure the length of the road doesn't get mistaken, six months in, for the road going nowhere. A nervous system that outsourced its own regulation to a substance every two to three hours is capable of running that regulation itself again. It just takes the time the biology actually requires, not the time anyone wishes it took.

END OF GUIDE — DISCONTINUANCE PROTOCOL: GHB / GBL / 1,4-BD

Harm Reduction Resource — Not Medical Advice — Not a Substitute for Emergency Care

Crisis Resources: 988 (Suicide & Crisis Lifeline, call or text) | Crisis Text: HOME to 741741 | SAMHSA: 1-800-662-4357